



PhytoTress™



Personalized Hair Care Solutions
With TrichoConcept™

Instructions For Use Booklet

AnazaoHealth® 
Corporation | Create. Heal. Live.®



This instructions for use booklet provides information about using PhytoTress™ a personalized hair care solution with TrichoConcept™, as well as certain active ingredient which may be prescribed by your physician.

Tell your physician and pharmacist about any drugs that you are taking (prescription or OTC, natural products, vitamins) and any health issues that you have. You must check to make sure that it is safe for you to take this drug with medication you may be taking and any health issues you may have.

Talk with your physician before starting any new drug, including prescription, OTC, natural product or vitamins.

None of the formulas described in this booklet are meant for infants or children unless specifically prescribed by a physician.

If you have additional questions, ask your physician or pharmacist.

Table of Contents

PhytoTress™	1
PhytoSolRx™	2,3
PhytoFoamRx™	4,5
PhytoCreamRx™	6
PhytoOilRx™	7
PhytoWashRx™	8
PhytoCondRx™	9
PhytoSerumRx™	10
Active Ingredients	11
Caffeine.....	12
Clobetasol.....	13,14
Finasteride.....	14,15
Ketoconazole.....	16,17
Latanoprost.....	18,19
Menthol.....	20
Minoxidil.....	21,22
Spirololactone.....	22,23
Vitamin A.....	24
Vitamin B.....	25,26
TrichoConcept™	27



General Information:

PhytoTress™ is a personalized hair care solution line with the patented TrichoTech™ Technology, acting on the balance of the entire hair system - bulb, scalp and shaft. It is 100% natural phytocomplex based on plant extracts. The patented TrichoTech™ Technology maximizes prescription results by the physiological improvement of the cellular environment. PhytoTress™ product line includes seven different vehicles that can be used individually or in combination with certain pharmaceutical ingredients for maximum results.

These seven vehicles are:

PhytoSolRx™	Non-irritating hydrophilic solution
PhytoFoamRx™	Foaming vehicle for localized application
PhytoCreamRx™	Ideal for eyelashes and eyebrows
PhytoOilRx™	Hydration from scalp to hair shaft
PhytoWashRx™	Cleans from root to tip without causing damage
PhytoCondRx™	Seals and strengthens hair cuticle
PhytoSerumRx™	Anti-frizz finishing



Non-irritating hydrophilic solution

PhytoSolRx™, compounded with TrichoSol™, is a highly spreadable hydrophilic solution containing mineral salts of vegetable origin. It is free from alcohol and propylene glycol. It is non-irritating and does not cause hair dryness. PhytoSolRx™ is an ideal and stable vehicle for a number of active pharmaceutical ingredients.

When using PhytoSolRx™ for the first time, remove the regular cap and replace it with the dropper cap.

Both caps are child resistant. To remove the cap, push down and turn in the direction of the arrows.

Directions For Use:

- Squeeze the rubber bulb and insert the dropper into the liquid. Release the bulb to fill the dropper to the one (1) mL line. Use only enough solution to cover the area to be treated. It is possible that it will be less than one (1) mL.
- Part your hair in the area to be treated. Place the tip of the dropper near the scalp and gently squeeze the bulb to gradually release the solution. Do this slowly and apply only a small amount at a time to prevent the solution from running off the scalp.
- Spread the solution evenly over the area to be treated. If you use your fingers, wash hands with soap and water immediately following application.
- After each use, attach the dropper cap to the bottle and close tightly.



PhytoSolRx™

Non-irritating hydrophilic solution

Other Important Information:

- PhytoSolRx™ is for external use only; specifically, the scalp. Do not use on other parts of the body.
- If your physician prescribed minoxidil to be added to PhytoSolRx™, do not use a hairdryer after applying. It may make the treatment less effective.
- Do not shampoo your hair for 4 hours after application if your physician prescribed minoxidil to be added to PhytoSolRx™.
- If you miss a dose, just continue with your next scheduled dose.
- Store at room temperature (68° to 77° F). Keep tightly closed.
- Allow the PhytoSolRx™ with minoxidil to completely dry for 2 to four hours after applying, including before going to bed. Minoxidil can stain clothing, hats, or bed linen if your hair or scalp is not fully dry after using this medicine.

Warnings:

- Avoid contact with the eyes. In case of accidental contact, rinse eyes with large amounts of cool, tap water.
- Do not use if your scalp is red, inflamed, infected, or painful.
- Contact your physician if scalp irritation or redness occurs.

Prescriptions are typically for a 30-day supply. Contact your physician if you are running out too soon.

Disclaimer: Some active ingredients require additional solvents.



PhytoFoamRx™

Foaming vehicle for localized application

PhytoFoamRx™, compounded with TrichoFoam™, is an alcohol-free and propylene glycol-free foam vehicle for water soluble and some liposoluble actives. Formulated with gentle, non-irritating and sensory-pleasing ingredients. PhytoFoamRx™ has been shown to be an ideal and stable vehicle for a number of active pharmaceutical ingredients. It is a soft foam with minimal oiliness and is free of propellants.

Directions For Use:

- Depress the pump once to dispense the foam into your palm.
- Apply to affected areas by massaging into the scalp. Do not rinse out.
- Wash hands immediately after use.

Other Important Information:

- PhytoFoamRx™ is for external use only; specifically, the scalp. Do not use on other parts of the body.
- If your physician prescribed minoxidil to be added to PhytoFoamRx™, do not use a hairdryer after applying. It may make the treatment less effective.
- Do not wash your hair for 4 hours after application if your physician prescribed minoxidil or to be added to PhytoFoamRx™.
- If you miss a dose, just continue with your next scheduled dose.
- Store at room temperature (68° to 77° F). Keep tightly closed.
- Allow the PhytoFoamRx™ with minoxidil to completely dry for 2-4 hours after applying, including before going to bed. Minoxidil can stain clothing, hats, or bed linen if your hair or scalp is not fully dry after using this medicine.



PhytoFoamRx™

Foaming vehicle for localized application

Warnings:

- Avoid contact with the eyes. In case of accidental contact, rinse eyes with large amounts of cool, tap water.
- Do not use if your scalp is red, inflamed, infected, or painful.
- Contact your physician if scalp irritation or redness occurs.

Prescriptions are typically for a 30-day supply. Contact your physician if you are running out too soon.

Disclaimer: Some active ingredients require additional solvents.



PhytoCreamRx™

Ideal for the eyelash and eyebrow area

PhytoCreamRx™, compounded with TrichoCream™, is a natural hydrophilic cream made from refined olive oil and shea butter that is. It is formulated with natural emulsifiers that contain high concentrations of triglycerides of linoleic and oleic acids. PhytoCreamRx™ is designed for use on eyelashes and eyebrows.

Directions For Use:

- Use the applicator brush to carefully apply the cream to the base of the eyelashes. Avoid contact with the eye.
- Use the applicator brush to apply enough cream to cover the eyebrow area.
- Discard applicator brush after each use.
- Wash hands after applying cream.

Other Important Information:

- PhytoCreamRx™ is for external use only.
- If you miss a dose, just continue with your next scheduled dose.
- Store at room temperature (68° to 77° F). Keep tightly closed.

Warnings:

- Avoid contact with the eyes. In case of accidental contact, rinse eyes with large amounts of cool, tap water.
- Do not use if the area being treated is red, inflamed, infected, or painful.
- Contact your physician if irritation or redness occurs at the treatment site.

Prescriptions are typically for a 30-day supply. Contact your physician if you are running out too soon.

Disclaimer: Some active ingredients require additional solvents.



PhytoOilRx™

Hydration from the scalp to the shaft

PhytoOilRx™, compounded with TrichoOil™, is a 100% natural lipid vehicle that contains essential fatty acids to promote restoration and hydration of the scalp and shaft and is. This product may be prescribed with additional vitamins as prescribed by your physician. It can be applied one or two times per week and is suitable for all hair types.

Directions For Use:

PhytoOilRx™ should be applied prewash, always on dry hair, and without any cosmetics applied.

- Place a small amount into the palm of the hand. The amount of oil needed will vary depending on the hair length.
- Proceed to rub hands together to coat them with oil on the palm.
- Beginning at the scalp, coat all your hair with the oil, working slowly from root to tip.
- Massage the oil over the entire scalp for up to three minutes.
- Repeat this process ensuring to work slowly from root to tip for best results.
- Rinse the hair liberally with water to remove excess oil before shampooing.

IMPORTANT: DO NOT LEAVE ON HAIR LONGER THAN TEN MINUTES! EXCESS OIL WILL BE DIFFICULT TO WASH OUT IF LEFT ON THE HAIR FOR AN EXTENDED PERIOD.

Prescriptions are typically for a 30-day supply. Contact your physician if you are running out too soon.

Disclaimer: Some active ingredients require additional solvents.



PhytoWashRx™

Clean from root to tip without causing damage

PhytoWashRx™, compounded with TrichoWash™, is a non-irritating shampoo that cleans the hair and scalp without destroying fatty acids. It is an extremely delicate, hydrophilic solution free of alcohol and propylene glycol. The PhytoWashRx™ formula promotes lasting hydration while adding softness and flexibility to the hair. With its potent antioxidant action, PhytoWashRx™ primes the hair for the delivery of selected pharmaceutical ingredients. The shampoo can be used on all types of hair.

Directions For Use:

- Wet the hair and scalp.
- Apply the same amount of shampoo as you normally use.
- Shampoo hair and gently massage the scalp.
- Allow the shampoo to remain for two minutes. Rinse and repeat.

When shampooing after applying PhytoOilRx™, it is important to rinse hair well with water to remove excess oil.

Prescriptions are typically for a 30-day supply. Contact your physician if you are running out too soon.

Disclaimer: Some active ingredients require additional solvents.



PhytoCondRx™

Sealing & strengthening the hair cuticle

PhytoCondRx™, compounded with TrichoCond™, is a high-power moisturizing conditioner, keeping the structure of hair aligned and firm. The adjusted pH of PhytoCondRx™ (always less than 4.5) promotes the final cohesion of the outer protective layer of the hair cell cuticles. PhytoCondRx™ provides a plentiful source of peptides needed to maintain the proteins already present in the hair cuticle. The peptides work in the hair cuticle by regulating the isoelectric point of the hair. This prevents cuticle breakage and exposure of the cortex. It increases lightness and vitality of the hair.

Directions for use:

- Wash hair with PhytoWashRx™, rinse, then remove excess water.
- Squeeze the bottle to dispense the conditioner into your hand. The amount needed to coat the hair will depend on hair length. If you have short hair, you will use a quarter size amount. If you have longer hair you will need more.
- Apply PhytoCondRx™ throughout the length of hair, from the scalp to the hair ends.
- Leave in the conditioner for about two minutes, then rinse.

For best results, use after PhytoWashRx™ and PhytoOilRx™

Prescriptions are typically for a 30-day supply. Contact your physician if you are running out too soon.

Disclaimer: Some active ingredients require additional solvents.



PhytoSerumRx™

Anti-frizz finishing serum

PhytoSerumRx™, compounded with TrichoSerum™, is a unique hair texturing system enhanced with patented smart TrichoTech™ Technology that is designed to provide: three-dimensional brightness, smoothness, and anti-frizz action. The serum improves the sealing of the hair cuticle while offering effective heat and color protection. For optimal use, it should be used on hair after conditioning with PhytoCondRx™. It can be used daily and does not create build up.

Directions For Use:

- Shampoo hair with PhytoWashRx™, condition hair with PhytoCondRx™ then rinse.
- Apply PhytoSerumRx™ to damp hair, and coat from root to tip.
- PhytoSerumRx™ is for the hair shaft only. There is no need to massage it into the scalp.
- Use a wide-tooth comb to detangle the hair if necessary.
- If you will be applying PhytoSolRx™ or PhytoFoamRx™, towel dry the hair first.

Prescriptions are typically for a 30-day supply. Contact your physician if you are running out too soon.

Disclaimer: Some active ingredients require additional solvents.

Active Ingredients

Caffeine

You should share the following information with your doctor before using this medication:

- If you have an allergy to caffeine, any other medications or food products. Be sure to explain to your doctor what symptoms you experienced, such as: rash, itching, shortness of breath, swelling, wheezing, cough.
- Ensure that your doctor is aware of all the medications (over the counter, prescription) you are currently taking as well as all medical conditions you have been diagnosed with.

Information to know while using this medication:

- Do not put on open wounds.
- It is for external use only. Do not take by mouth.
- Wash hands after use.
- Follow the directions as prescribed by the doctor. Do not use more often than directed.

Contact your doctor immediately if you experience:

- Signs of allergic reaction, like hives, rash, itching, red, swollen, or blistered skin with or without fever, wheezing, tightness in the chest, trouble breathing, swallowing or talking, hoarseness, swelling of the mouth, face, lips, tongue or throat.

What to do if you miss a dose:

- Take the next scheduled dose.

Storage and other information:

- Store at room temperature.
- Store all medications in a secure area out of reach of children and pets.

Clobetasol

Pregnancy: Clobetasol is classified as FDA pregnancy risk category C. Although there are no adequate and well-controlled studies in pregnant women, limited experience in human pregnancy has not resulted in clinically significant risk to the fetus.

You should share the following information with your doctor before using this medication:

- If you have an allergy to Clobetasol, any medications or food products. Be sure to explain to your doctor what symptoms you experienced, such as: rash, itching, shortness of breath, swelling, wheezing, cough.
- If you have thinning of the skin where you will be applying this medication.
- If you have an infection in the area you will be applying this medication.
- Ensure that your doctor is aware of all the medications (over the counter, prescription) you are currently taking as well as all medical conditions you have been diagnosed with.
- Speak with your doctor if you will be applying other medications to the treatment area(s).
- Tell your doctor if you are pregnant or plan on getting pregnant. You will need to talk about the benefits and risks of using this drug while you are pregnant.
- Tell your doctor if you are breast-feeding. You will need to talk about any risks to your baby.

Information to know while using medication:

- Do not apply on cut or damaged skin.
- Follow the directions as prescribed by the doctor. Do not use more often than directed.
- For external use only. Do not take by mouth.
- Do not cover head while shampoo is on the head.
- Do not put shampoo on the face, underarms or groin area.

Contact your doctor immediately if you experience:

- Signs of allergic reaction such as: hives, rash, itching, red, swollen, blistered skin with or without fever, wheezing, tightness in the chest, trouble breathing, swallowing or talking, hoarseness, swelling of the mouth, face, lips, tongue or throat.

Clobetasol

- Signs of elevated blood sugar such as: confusion, sleepiness, thirstiness, hunger, increased urination, flushing, fast breathing, or fruity breath smell.
- Signs of Cushing's disease such as: weight gain in the upper back or belly, moon face, severe headache, or slow healing.
- Signs of a weak adrenal gland such as: a very bad upset stomach or throwing up, very bad dizziness or passing out, muscle weakness, feeling very tired, mood changes, a decrease in appetite, or weight loss.
- Irritation where this medication is used.
- Thinning of the skin.
- Change in eyesight, eye pain, or very bad eye irritation.

Other potential side effects:

- Stinging, burning, redness.
- These are not the only possible side effects. If you have questions about side effects, contact your physician or pharmacist.
- May increase risk of forming a cataract or may increase the intraocular pressure of the eye leading to glaucoma.

Storage and other information:

- Store at room temperature.
- Store all medications in a secure area out of reach of children and pets.

Finasteride

This medication is NOT approved for women. If you are a woman considering using this medication, please speak with your doctor.

You should share the following information with your doctor before using this medication:

- If you have an allergy to Finasteride, any medications or food products. Be sure to explain to your doctor what symptoms you experienced, such as: rash, itching, shortness of breath, swelling, wheezing, cough.
- Ensure that your doctor is aware of all the medications (over the counter, prescription) you are currently taking as well as all medical conditions you have been diagnosed with.

Finasteride

Information to know while using medication:

- Women should not handle finasteride. If a woman is to come in contact with the medication, wash the exposed area thoroughly.
- Use it on scalp and hair only. Do not take by mouth.
- Avoid driving and doing other tasks or actions that call for you to be alert until you see how this medication affects you.
- To lower the chance of feeling dizzy or passing out, rise slowly if you have been sitting or lying down. Be careful going up and down stairs.
- This drug may affect certain lab tests. Be sure to inform your health care providers that you are currently undergoing therapy with this medication.
- Do not donate blood while using this medication and for 1 month after discontinuing treatment.
- Follow the directions as prescribed by your doctor. Do not use more often than directed.
- This medication may lower sperm count. Ask your doctor if this medication would hinder the chances of fathering a child.

Contact your doctor immediately if you experience:

- Signs of allergic reaction, like hives, rash, itching, red, swollen, or blistered skin with or without fever, wheezing, tightness in the chest, trouble breathing, swallowing or talking, hoarseness, swelling of the mouth, face, lips, tongue or throat.
- Very bad dizziness or passing out.
- Lump in the breast, breast soreness
- Testicular pain

Other potential side effects:

- Dizziness, feeling tired or weak.
- May cause a change in sex ability in men, or lowered sex interest.
- These are not the only potential side effects. If you have questions about side effects, contact your physician.

What to do if you miss a dose:

- Take your next scheduled dose.

Storage and other information:

- Store at room temperature. Store in a dry place.
- Store all medications in a secure area out of reach of children and pets.

Ketoconazole

Pregnancy: Ketoconazole is classified as FDA pregnancy risk category C. Although there are no adequate and well-controlled studies in pregnant women, limited experience in human pregnancy has not resulted in clinically significant risk to the fetus.

You should share the following information with your doctor before using this medication:

- If you have an allergy to Ketoconazole, any medications or food products. Be sure to explain to your doctor what symptoms you experienced, such as: rash, itching, shortness of breath, swelling, wheezing, cough.
- Ensure that your doctor is aware of all the medications (over the counter, prescription) you are currently taking as well as all medical conditions you have been diagnosed with.
- Tell your doctor if you are breast-feeding. You will need to discuss risks to you and your baby.

Information to know while using medication:

- Follow the directions as prescribed by the doctor. Do not use more often than directed.
- For external use only. Do not take by mouth.
- Use shampoo on hair and scalp only. Keep out of mouth, nose ears and eyes.

What to do if you miss a dose:

- Take the next scheduled dose.

Contact your doctor immediately if you experience:

- Signs of allergic reaction such as: hives, rash, itching, red, swollen, blistered skin with or without fever, wheezing, tightness in the chest, trouble breathing, swallowing or talking, hoarseness, swelling of the mouth, face, lips, tongue or throat.
- Very bad irritation where the medication is used.

Ketoconazole

Other potential side effects:

- Burning or irritation where the medication is used.
- May take curl out of hair.
- Change in hair color or texture
- Itching, dry skin.
- These are not the only possible side effects. If you have questions about side effects, contact your physician.

Storage and other information:

- Store at room temperature.
- Protect from light.
- Store all medications in a secure area out of reach of children and pets.

Latanoprost

Pregnancy: Latanoprost is classified as FDA pregnancy risk category C. Although there are no adequate and well-controlled studies in pregnant women, limited experience in human pregnancy has not resulted in clinically significant risk to the fetus.

Breast Feeding: It is not known whether latanoprost or its metabolites are excreted in breast milk. Consider the benefits of breast-feeding and risk of potential infant drug exposure versus an untreated or treated condition.

You should share the following information with your doctor before using this medication:

- If you have an allergy to Latanoprost, any medications or food products. Be sure to explain to your doctor what symptoms you experienced, such as: rash, itching, shortness of breath, swelling, wheezing, cough.
- Ensure that your doctor is aware of all the medications (over the counter, prescription) you are currently taking as well as all medical conditions you have been diagnosed with.
- If you are pregnant or plan to become pregnant.

Information to know while using medication:

- Follow the directions as prescribed by the doctor. Do not use more often than directed.
- Do not apply to red, swollen, irritated or painful skin.
- When applying to eyelashes, always use a new applicator brush.
- Do not use other scalp drugs while using Latanoprost/Minoxidil in PhytoTress™ bases.
- It may take several months to see the full effect.
- For use on scalp and hair only; however, PhytoCream™ with Latanoprost may be used on eyelashes and eyebrows.
- Do not take by mouth.
- Wash hands thoroughly after use.
- If this medication gets into the eyes, rinse with cool tap water.
- Do not use within 24 hours of hair coloring treatment.

Latanoprost

Contact your doctor immediately if you experience:

- Signs of allergic reaction, like hives, rash, itching, red, swollen, or blistered skin with or without fever, wheezing, tightness in the chest, trouble breathing, swallowing or talking, hoarseness, swelling of the mouth, face, lips, tongue or throat.

Other potential side effects:

- Eye irritation
- This is not the only possible side effect. If you have questions about side effects, contact your physician.

What to do if you miss a dose:

- Take your next scheduled dose.

Storage and other information:

- Store at room temperature. Store in a dry place.
- Store all medications in a secure area out of reach of children and pets.

Menthol

You should share the following information with your doctor before using this medication:

- If you have an allergy to menthol, any other medications or food products. Be sure to explain to your doctor what symptoms you experienced, such as: rash, itching, shortness of breath, swelling, wheezing, cough.
- Ensure that your doctor is aware of all the medications (over the counter, prescription) you are currently taking as well as all medical conditions you have been diagnosed with.

Information to know while using this medication:

- Do not put on open wounds.
- It is for external use only. Do not take by mouth.
- Wash hands after use.
- Follow the directions as prescribed by the doctor. Do not use more often than directed.

Contact your doctor immediately if you experience:

- Signs of allergic reaction, like hives, rash, itching, red, swollen, or blistered skin with or without fever, wheezing, tightness in the chest, trouble breathing, swallowing or talking, hoarseness, swelling of the mouth, face, lips, tongue or throat.

What to do if you miss a dose:

- Take your next scheduled dose.

Storage and other information:

- Store at room temperature. Protect from heat.
- Store all medications in a secure area out of reach of children and pets.

Minoxidil

Pregnancy: Minoxidil is classified as FDA pregnancy risk category C. Although there are no adequate and well-controlled studies in pregnant women, limited experience in human pregnancy has not resulted in clinically significant risk to the fetus.

You should share the following information with your doctor before using this medication:

- If you have an allergy to minoxidil, any other medications or food products. Be sure to explain to your doctor what symptoms you experienced, such as: rash, itching, shortness of breath, swelling, wheezing, cough.
- If you do not have a family history of hair loss or if your hair loss was sudden.
- Ensure that your doctor is aware of all the medications (over the counter, prescription) you are currently taking as well as all medical conditions you have been diagnosed with.
- If you are pregnant or plan to become pregnant.

Information to know while using medication:

- Follow the directions as prescribed by the doctor. Do not use more than prescribed.
- Do not apply to red, swollen, irritated or painful skin.
- Do not use other scalp drugs while using Minoxidil/Latanoprost in PhytoTress™ base.
- It may take several months to see the full effect.
- For use on scalp and hair only. Do not take by mouth.
- Wash hands thoroughly after use.
- If this drug gets into the eyes, rinse with cool tap water.
- Do not use within 24 hours of hair coloring treatment.

Contact your doctor immediately if:

- You show signs of allergic reaction. Such as: hives, rash, itching, red, swollen, or blistered skin with or without fever, wheezing, tightness in the chest, trouble breathing, swallowing or talking, hoarseness, swelling of the mouth, face, lips, tongue or throat.
- Chest pain, pressure or fast heartbeat.
- Swelling in the arms or legs.
- Scalp irritation.

Minoxidil

Other potential side effects:

- Change in hair color or texture.
- This is not the only possible side effect. If you have questions about side effects, contact your physician.

What to do if you miss a dose:

- Take the next scheduled dose.

Storage and other information:

- Store at room temperature. Store in a dry place.
- Store all medications in a secure area out of reach of children and pets.

Spironolactone

Pregnancy: Spironolactone is classified as FDA pregnancy risk category C. Although there are no adequate and well-controlled studies in pregnant women, limited experience in human pregnancy has not resulted in clinically significant risk to the fetus.

The information you should share with your doctor before using this medication:

- If you have an allergy to Spironolactone, any medications or food products. Be sure to explain to your doctor what symptoms you experienced, such as: rash, itching, shortness of breath, swelling, wheezing, cough.
- If you are pregnant or plan to get pregnant. You will need to talk about the benefits and risks of using this medication while you are pregnant.
- If you have Addison's disease.
- If you have high potassium levels.
- If you are taking amiloride, eplerenone, or triamterene.
- If you are on a low salt diet.
- If you have diabetes.
- If you have kidney disease.
- Ensure that your doctor is aware of all the medications (over the counter, prescription) you are currently taking as well as all medical conditions you have been diagnosed with.

Spironolactone

Information to know while using this medication:

- Avoid driving and other tasks or actions that call for you to be alert until you see how this medication affects you.
- This medication may affect certain lab tests. Tell all health care providers and lab workers you take this medication.
- Follow the directions as prescribed by the doctor. Do not use more often than directed.

Contact your doctor if:

- Signs of allergic reaction, like hives, rash, itching, red, swollen, or blistered skin with or without fever, wheezing, tightness in the chest, trouble breathing, swallowing or talking, hoarseness, swelling of the mouth, face, lips, tongue or throat.
- Signs of mood changes, confusion, muscle pain or weakness, a heartbeat that does not feel normal, very bad dizziness, passing out, excessive thirst, change in urine output, very bad upset stomach, or throwing up.
- Lowered interest in sex, not able to get or keep an erection.
- Fever, chills, sore throat.
- Unexplained bruising or bleeding.
- Breast pain, menstrual changes.

Other potential side effects:

- Loose stools, diarrhea.
- Sleepiness, dizziness, headache.
- Upset stomach.
- These are not the only possible side effect. If you have questions about side effects, contact your physician.

What to do if you miss a dose:

- Take the next scheduled dose.

Storage and other information:

- Store at room temperature.
- Store all medications in a secure area out of reach of children and pets.

Vitamin A

The information you should share with your doctor before using this medication:

- If you have an allergy to vitamin A, any other medications or food products. Be sure to explain to your doctor what symptoms you experienced, such as: rash, itching, shortness of breath, swelling, wheezing, cough.
- Ensure that your doctor is aware of all the medications (over the counter, prescription) you are currently taking as well as all medical conditions you have been diagnosed with.
- Information to know while using this medication:
- Do not put on open wounds.
- It is for external use only. Do not take by mouth.
- Wash hands after use.
- Follow the directions as prescribed by the doctor. Do not use more often than directed.

Contact your doctor immediately if you experience:

- Signs of allergic reaction, like hives, rash, itching, red, swollen, or blistered skin with or without fever, wheezing, tightness in the chest, trouble breathing, swallowing or talking, hoarseness, swelling of the mouth, face, lips, tongue or throat.

What to do if you miss a dose:

- Take the next scheduled dose.

Storage and other information:

- Store at room temperature.
- Store all medications in a secure area out of reach of children and pets.

Vitamin B6

Pregnancy: Vitamin B is classified as FDA pregnancy risk category A. Although there are no adequate and well-controlled studies in pregnant women, limited experience in human pregnancy has not resulted in clinically significant risk to the fetus.

You should share the following information with your doctor before using this medication:

- If you have an allergy to Vitamin B, any medications or food products. Be sure to explain to your doctor what symptoms you experienced, such as: rash, itching, shortness of breath, swelling, wheezing, cough.
- If you are pregnant or plan to get pregnant. You will need to talk about the benefits and risks of using this medication while you are pregnant.
- Tell your doctor if you are breast-feeding. You will need to discuss risks to your baby.
- Ensure that your doctor is aware of all the medications (over the counter, prescription) you are currently taking as well as all medical conditions you have been diagnosed with.

Information to know while using medication:

- Follow the directions as prescribed by the doctor. Do not use more often than directed.
- Contact your doctor immediately if you experience: signs of allergic reaction, like hives, rash, itching, red, swollen, or blistered skin with or without fever, wheezing, tightness in the chest, trouble breathing, swallowing or talking, hoarseness, swelling of the mouth, face, lips, tongue or throat.

Other side effects:

- A burning, numbness, or tingling sensation that is not normal.
- This is not the only possible side effect. If you have questions about side effects, contact your physician.

Vitamin B6

What to do if you miss a dose:

- Take your next scheduled dose.

Storage and other information:

- Store at room temperature.
- Store all medications in a secure area out of reach of children and pets.

Disclaimer: The headings and descriptions contained within this document are general headings describing formulas, products and services offered by AnazaoHealth Corporation. These general headings are not intended and should not be interpreted to make recommendations or claims regarding the use, efficacy or safety of products, formulas and/or services listed herein. Only a doctor or other appropriately licensed health care professional, as a learned intermediary, can determine if a formula, product or service described herein is appropriate for a patient.

TrichoConcept™

7 vehicles with Patented TrichoTech™ Technology



The first global line of multifunctional vehicles with TrichoTech™ Technology, that balances the entire hair system - bulb, scalp and shaft.



TrichoSol™



TrichoFoam™



TrichoCream™



TrichoOil™



TrichoWash™



TrichoCond™



TrichoSerum™

TrichoConcept™* is free of

Parabens, Formaldehyde, P-Amino Benzoic Acid, Boric Acid, 1,4-Dioxane, Benzyl Alcohol Benzyl Benzoate, Propylene Glycol, Triclosan, Peanut Oil, Mineral Oil Petrolatum, Lanolin, and Artificial Colorants.

**TrichoConcept™ bases by Fagron, Inc.*

Contact Us

Shipping Address:

AnazaoHealth Corporation
5710 Hoover Boulevard
Tampa, Florida 33634

Phone: 800.995.4363

Hours of Operation:

Monday-Friday
8:00 a.m. - 8:30 p.m. EST

Customer Service Email:

customerservice@anazaohealth.com

Online Ordering Website:

www.myAnazao.com

Website:

www.anazaohealth.com

The trademarks listed herein are the property of their respective owners. AnazaoHealth®, Anazao®, Create, Heal, Live.®, and the Compass are registered trademarks of AnazaoHealth Corporation, © 2019. All Rights Reserved. APT601-032619